VOL. 595, NOS. 1+2 JOURNAL OF CHROMATOGRAPHY MARCH 20, 1992

CONTENTS

(Abstracts/Contents Lists published in Analytical Abstracts, Biochemical Abstracts, Biological Abstracts, Chemical Abstracts, Chemical Titles, Chromatography Abstracts, Current Contents/Life Sciences, Current Contents/Physical, Chemical & Earth Sciences, Deep-Sea Research/Part B: Oceanographic Literature Review, Excerpta Medica, Index Medicus, Mass Spectrometry Bulletin, PASCAL-CRNS, Referativnyi Zhurnal, Research Alert and Science Citation Index)

REVIEW

Polychlorinated biphenyls in the environment
V. Lang (Brno, Czechoslovakia) 1

REGULAR PAPERS

Column Liquid Chromatography

Solvatochromic hydrogen bond donor acidity of cyclodextrins and reversed-phase liquid chromatographic retention of small molecules on a β-cyclodextrin-bonded silica stationary phase
by J. H. Park, M. D. Jang and M. J. Shin (Kyongsan, South Korea) (Received October 30th, 1991) 45

Validation of chromatographic retention models in reversed-phase high-performance liquid chromatography by fitting experimental data to the relevant equations
by N. Sadlej-Sosnowska and I. Śledzińska (Warsaw, Poland) (Received October 8th, 1991) 53

Chromatographic resolution on methylbenzoylcellulose beads. Modulation of the chiral recognition by variation of the position of the methyl group on the aromatic ring
by E. Francotte and R. M. Wolf (Basle, Switzerland) (Received December 5th, 1991) 63

Preparation of nitrophenylethylsilylated silica gel and its chromatographic properties in the separation of polychlorinated dibenzo-*p*-dioxins
by K. Kimata, K. Hosoya, N. Tanaka and T. Araki (Kyoto, Japan) and D. G. Patterson, Jr. (Atlanta, GA, USA) (Received November 25th, 1991) 77

Evaluation of reversed-phase liquid chromatographic columns for recovery and selectivity of selected carotenoids
by K. S. Epler and L. C. Sander (Gaithersburg, MD, USA), R. G. Ziegler (Bethesda, MD, USA) and S. A. Wise and N. E. Craft (Gaithersburg, MD, USA) (Received November 28th, 1991) 89

Physical and biochemical characterization of five commercial resins for immunoaffinity purification of factor IX
by J. Tharakan (Washington, DC, USA), F. Highsmith (College Park, MD, USA) and D. Clark and W. Drohan (Rockville, MD, USA) (Received November 19th, 1991) 103

Immobilization of monoclonal antibodies for affinity chromatography using a chelating peptide
by P. Loetscher, L. Mottlau and E. Hochuli (Basle, Switzerland) (Received November 26th, 1991) 113

Stationary phase effect on enantioselectivity of dansyl phenylalanine in microcolumn liquid chromatography with γ-cyclodextrin as mobile phase additive
by T. Takeuchi (Nagoya, Japan) and N. Nagae (Seto, Japan) (Received November 21st, 1991) 121

Reversed-phase chromatographic behaviour of β-endorphin: evidence of conformational change
by L. S. Monger and C. J. Olliff (Brighton, UK) (Received October 29th, 1991) 125

Comparison of colorimetric and chromatographic determination of *tert.*-butylpyrocatechol
by E. Hlavačková and R. Reya (Bratislava, Czechoslovakia) (Received November 12th, 1991) 137

Fluorescence detection of trichothecene mycotoxins as coumarin-3-carbonyl chloride derivatives by high-performance liquid chromatography
by H. Cohen and B. Boutin-Muma (Ottawa, Canada) (Received November 28th, 1991) 143

Ion-exclusion chromatography using mobile phases containing β-cyclodextrin
by B. K. Głod, P. R. Haddad and P. W. Alexander (Kensington, Australia) (Received November 20th, 1991) 149

Quantitative analysis of total resin acids by high-performance liquid chromatography of their coumarin ester derivatives
by D. E. Richardson (Boyer, Australia) and J. B. Bremner and B. V. O'Grady (Hobart, Australia) (Received December 3rd, 1991) 155

(Continued overleaf)

Contents (continued)

High-performance liquid chromatography of histamine and 1-methylhistamine with on-column fluorescence derivatization
by K. Saito, M. Horie and N. Nose (Saitama, Japan), K. Nakagomi (Ibaragi, Japan) and H. Nakazawa (Tokyo, Japan) (Received November 11th, 1991) 163

Purity assay of sodium mercaptododecarborate by high-performance liquid chromatography
by B. Grüner and Z. Plzák (Řež, Czechoslovakia) and I. Vinš (Prague, Czechoslovakia) (Received November 25th, 1991) 169

High-performance liquid chromatography of sulphonamides extracted from bovine and porcine muscle by solid-phase dispersion
by L. V. Walker, J. R. Walsh and J. J. Webber (Hamilton, Australia) (Received December 16th, 1991) 179

Aqueous high-performance size-exclusion chromatographic assay for high-molecular-weight impurities in ceftiofur sodium
by M. J. Dunn and D. A. Hahn (Kalamazoo, MI, USA) (Received November 4th, 1991) 185

Screening of organophosphorus pesticides using liquid chromatography–atmospheric pressure chemical ionization mass spectrometry
by S. Kawasaki and H. Ueda (Saga, Japan), H. Itoh (Tokyo, Japan) and J. Tadano (Saga, Japan) (Received October 28th, 1991) 193

Stereochemical analysis of betamethasone and dexamethasone by derivatization and high-performance liquid chromatography
by S.-H. Chen, S.-M. Wu and H.-L. Wu (Kaohsiung, Taiwan) (Received November 12th, 1991) 203

Gas Chromatography

Critical capillary column examination of the relationship between separation number and height equivalent to a theoretical plate and their dependence on temperature
by L. A. Jones, J. J. Glennon and W. H. Reiss (Raleigh, NC, USA) (Received August 20th, 1991) 209

Van Deemter-type relationship for determining the optimum initial flow-rate and optimum pressure programming rate in temperature/pressure-programmed capillary column gas chromatography utilizing separation numbers
by L. A. Jones, W. H. Reiss, J. J. Glennon and T. M. Gerig (Raleigh, NC, USA) (Received August 20th, 1991) . . . 221

Activity coefficients in binary liquid mixtures measured by reversed-flow gas chromatography
by A. Koliadima, G. Karaiskakis and N. A. Katsanos (Patras, Greece) and M. Roth (Brno, Czechoslovakia) (Received November 20th, 1991) 237

Gas chromatographic analysis of high-molecular-weight polynuclear aromatic hydrocarbons. I. Molecular weight 328
by A. Bemgård, A. Colmsjö and B.-O. Lundmark (Solna, Sweden) (Received November 27th, 1991) 247

Determination of α- and β-zearalenol and zearalenone in cereals by gas chromatography with ion-trap detection
by K. Schwadorf and H.-M. Müller (Stuttgart, Germany) (Received September 5th, 1991) 259

Analytical methods for monoterpene glycosides in grape and wine. II. Qualitative and quantitative determination of monoterpene glycosides in grape
by S. G. Voirin, R. L. Baumes, J.-C. Sapis and C. L. Bayonove (Montpellier, France) (Received November 5th, 1991) 269

Degradation of polysulphones by alkali fusion
by J. K. Haken and M. Camamo (Kensington, Australia) (Received November 14th, 1991) 283

Automated clean-up procedure for the gas chromatographic–high-resolution mass spectrometric determination of polychlorinated dibenzo-*p*-dioxins and dibenzofurans in milk
by J. A. van Rhijn, W. A. Traag, W. Kulik and L. G. M. Th. Tuinstra (Wageningen, Netherlands) (Received November 21st, 1991) 289

Supercritical Fluid Chromatogaphy

Quantitative hydrocarbon group analysis of gasoline and diesel fuel by supercritical fluid chromatography
by P. E. Andersson, M. Demirbüker and L. G. Blomberg (Stockholm, Sweden) (Received November 20th, 1991) . . 301

Electrophoresis

Peak dispersion and separation efficiency in high-performance zone electrophoresis with gel-filled capillaries
by E. Kenndler and C. Schwer (Vienna, Austria) (Received November 13th, 1991) 313

High-sensitivity laser-induced fluorescence detection of native proteins in capillary electrophoresis
by T. T. Lee and E. S. Yeung (Ames, IA, USA) (Received December 3rd, 1991) 319

Relationship between zone length and step height in isotachophoresis
by M. T. Ackermans, F. M. Everaerts and J. L. Beckers (Eindhoven, Netherlands) (Received October 11th, 1991) . . 327

SHORT COMMUNICATIONS

Column Liquid Chromatography

Chromatographic behaviour of dipyridylsulphides. Relationship between log k' values and structure by reversed-phase high-performance liquid chromatography
by J. Klimeš, V. Klimešová and K. Waisser (Hradec Králové, Czechoslovakia) (Received November 19th, 1991) . . . 334

High-performance liquid chromatography of isopropylphenols
by M. Petro, K. Križanovič and K. Ďulák (Bratislava, Czechoslovakia) (Received December 12th, 1991) 337

Analyses of isomeric mono-O-methyl-D-glucoses, D-glucobioses and D-glucose monophosphates by high-performance anion-exchange chromatography with pulsed amperometric detection
by K. Koizumi, Y. Kubota, H. Ozaki, K. Shigenobu, M. Fukuda and T. Tanimoto (Nishinomiya, Japan) (Received December 13th, 1991) 340

Determination of maleic hydrazide in tobacco by micellar liquid chromatography
by S. S. Yang (Richmond, VA, USA) (Received December 18th, 1991) 346

Endogenous alkaloids in man. XI. Analysis of glyoxylate-derived 1,3-thiazolidines by ion-pair-assisted reversed-phase chromatography
by G. Bringmann, C. Hesselmann and D. Feineis (Würzburg, Germany) (Received December 16th, 1991) 351

Separation and purification of azetidinyl methyl sulfinates using preparative high-performance liquid chromatography
by F. Brown, Jr. and P. C. Heath (Indianapolis, IN, USA) (Received December 30th, 1991) 355

Gas Chromatography

Wide dynamic range electron-capture detection using the electron-capture detector pulse frequency signal
by R. J. Lagomarsino and N. Latner (New York, NY, USA) (Received December 18th, 1991) 359

Identification of ochratoxin A in food samples by chemical derivatization and gas chromatography–mass spectrometry
by Y. Jiao, W. Blaas, C. Rühl and R. Weber (Berlin, Germany) (Received December 9th, 1991) 364

Planar Chromatography

Comparison of commercially available thin-layer chromatography plates with mixtures of dyes, analgesics and phenols
by D. L. Gumprecht (Tuscaloosa, AL, USA) (Received January 3rd, 1992) 368

BOOK REVIEW

Chiral separations by liquid chromatography (edited by S. Ahuja), reviewed by W. Lindner 375

Author Index 377

Errata 379

Announcement of International Ion Chromatography Symposium 1992 380

In articles with more than one author, the name of the author to whom correspondence should be addressed is indicated in the article heading by a 6-pointed asterisk (*).